arXiv:physics/0104038v1 [physics.atom-ph] 10 Apr 2001

Cooling atoms in an optical trap by selective parametric excitation

N. Poli, R.J. Brecha*, G. Roati†, and G. Modugno
INFM, LENS, and Dipartimento di Fisica, Universitá di Firenze, Largo E. Fermi 2, 50125 Firenze, Italy

(February 2, 2008)

We demonstrate the possibility of energy-selective removal of cold atoms from a tight optical trap by means of parametric excitation of the trap vibrational modes. Taking advantage of the anharmonicity of the trap potential, we selectively remove the most energetic trapped atoms or excite those at the bottom of the trap by tuning the parametric modulation frequency. This process, which had been previously identified as a possible source of heating, also appears to be a robust way for forcing evaporative cooling in anharmonic traps.

32.80.Pj, 32.80.Lg

Parametric excitation of atomic motion due to laser intensity fluctuations has been recently indicated as one of the major sources of heating in far-off resonance optical traps (FORTs) [1,2], and as a limitation in achieving very long trapping times [3,4]. In this Letter we show that parametric excitation in a FORT can also lead to a cooling of the trapped sample, since the anharmonicity of the potential allows a selective removal of the most energetic atoms alone.

The excitation process has typically been modeled assuming a harmonic FORT potential [1,2]. For a modulation of the potential depth at twice the trap vibrational frequencies, an exponential heating of the trapped sample, accompanied by trap losses, has been predicted. The main features of the parametric heating process predicted by the harmonic theory have been experimentally observed [5–7]. Going beyond the basic trap model, it has been possible to further refine theoretical calculations through modeling of the anharmonic trapping potential used in actual experiments [8]. One of the consequences of the anharmonicity is a variation of the trap frequencies with vibrational energy, which implies that a modulation at a single frequency will be resonant with a few trap levels only.

We present here a detailed study of the temperature of a sample of cold atoms in a one-dimensional (1D) lattice FORT, in which the anharmonicity of the potential is exploited to perform energy-selective excitation and removal of the atoms, resulting in either heating or cooling of the remaining trapped sample. In particular, the cooling process appears to be a robust and relatively efficient way to lower the atomic temperature via forced evaporation.

The trap is created with a beam of single-mode radiation generated by a Ti:sapphire laser at λ_T=787 nm, red-detuned by approximately 18 nm from the D-lines of potassium. The laser beam, after passing through an acousto-optic modulator (AOM), is focused to w=90 μm within a two-lens telescope, and then retroreflected to form a vertical standing-wave. The focusing is chosen to yield a large Rayleigh length, z_R=3 cm, which results in an almost uniform lattice depth along the vertical direction. Therefore, the optical potential close to the beam waist has the form

$$U(r,z) = -U_0 \cos^2(kz) \exp\Big(-\frac{2r^2}{w^2}\Big), \qquad (0.1)$$

where $k = 2\pi/\lambda_T$. About 10^7 atoms are prepared in a magneto-optical trap (MOT) at a temperature of 60 μK, and typically 5% of them are transferred to the FORT thanks to a compression procedure described in [9]. After loading the atoms in the FORT, the MOT beams are extinguished by means of AOMs and mechanical shutters, and the atoms are left to evolve in the presence of the trap light only. The density and the radial temperature of the trapped atoms are determined using absorption imaging of the atomic cloud after 1 ms of expansion following release from the FORT. The typical vertical size of the cloud of 700 μm indicates that approximately 1800 lattice sites are occupied by an average of 300 atoms each. The axial temperature of the atoms is measured with a time-of-flight (TOF) technique, by absorption of a sheet of light passing about 1.5 cm below the FORT. The temperature of the trapped atoms, after a 100-ms equilibration phase following loading, stays constant during storage in the FORT [9], and the radial and axial temperatures are found to be equal within the uncertainty. The lifetime of the FORT is limited to about 1.2 s mainly by collisions with background gas.

The harmonic approximations for the radial and axial frequencies of the single lattice site are measured with a parametric excitation technique, as described below. Typical values for these quantities are ω_r=$2\pi\times$1.3 kHz, ω_a=$2\pi\times$700 kHz, which for the typical temperature T=80 μK lead to a density of 10^{12} cm^{-3}. From the measured value of ω_a it is possible to estimate the trap depth as $U_0 = M\lambda_T^2\omega_a^2/8\pi^2$, which corresponds to U_0=700 μK.

We apply the parametric excitation via the control AOM as a small sinusoidal modulation on the power of the trapping laser, typically 200 ms after loading the

---

[*] permanent address, Physics Dept., University of Dayton, Dayton, OH.

[†] also at Dipartimento di Fisica, Università di Trento, I-38050 Povo (Tn), Italy.

FORT. To measure the spectrum of excitation we have repeated at different modulation frequencies a procedure composed of an excitation for a period T_{ex} at fixed modulation amplitude ϵ and frequency Ω, followed by detection of the number of atoms and radial temperature. For both radial and axial excitation, as shown in Fig. 1 and Fig. 2 respectively, we see large resonances in the loss of atoms, accompanied by dispersive resonances in the temperature of the atoms left in the trap. The resonances in the trap losses have been observed previously in optical lattices [5,8,9], and their presence is expected even if we approximate the FORT as a harmonic potential: when Ω is close to $2\,\omega_a$ or $2\,\omega_r$ (or to their subharmonics) the atoms are excited upwards along the ladder of energy levels [1], and eventually lost from the trap. Note that for the radial excitation two resonances at about ω_r and $2\,\omega_r$ are present, as expected for a harmonic trap, while in the axial case there is an additional resonance close to $3.6\,\omega_a$, which is a clear signature of the anharmonicity of the trap [8].

On the other hand, the observed shape of the resonances in the temperature of the remaining trapped atoms are not expected for a harmonic trap [1,2]. Their origin must be related to the anharmonicity of the potential, which implies that the vibrational frequency changes with atomic energy. In Fig. 3 we show the excitation frequency $2\,\omega_a$ calculated for a perfectly 1D sinusoidal lattice: while the atoms at the bottom of the trap are expected to oscillate at the harmonic vibrational frequency, the most energetic ones have a much lower frequency, because of the decreased curvature of the potential. A similar behavior is also expected for the radial oscillation frequency because of the gaussian shape of the radial potential. Therefore, when the modulation frequency is tuned to the red of $2\,\omega_r$, $2\,\omega_a$ or their harmonics and subharmonics, it is mainly the most energetic atoms that are parametrically excited and ejected from the trap, resulting in a reduction of the mean energy of the remaining atoms. On the contrary, resonant modulation frequencies excite primarily atoms at the bottom of the trap, resulting in a net increase of the mean energy. We therefore identify the quantities $2\,\omega_r$, $2\,\omega_a$ with the position of the main resonances in the temperature spectra of Fig. 1 and Fig. 2 leading to a heating of the trapped sample.

These observations are an evidence of theoretical predictions that the peak in the resonance of atomic losses in the trap occurs at a slightly lower frequency than the actual harmonic frequencies [8]. This kind of information is very useful, since the frequencies measured from the trap loss resonances are used to estimate quantities such as the number density or the phase-space density of atoms in an optical lattice [5,9].

Note that the temperature measurements in Fig. 1 and Fig. 2 have been performed on the radial degree of freedom only, since the TOF detection of the axial temperature is strongly perturbed by the atoms excited out of the trap as long as 100 ms before release. In all the measurement we report here we have extracted the radial temperature from the width of the atomic distribution, which was always well fitted by a gaussian, assuming thermal equilibrium.

In the following we restrict our attention to the excitation of the axial vibrational mode, which is the most energetic and therefore requires a smaller perturbation of the trapping potential [10], making modeling easier. We also suspect that the large broadening of the radial resonances in Fig. 1 might come from an ellipticity of the laser beam that makes ω_r vary in the radial plane, masking interesting characteristics of the resonances' shape.

To analyze in more detail the axial resonances we need to refine the simple 1D model for the dispersion of the axial vibrational frequencies presented in Fig. 3. Indeed, when considering the actual potential of Eq. 0.1 one would expect that the axial and radial degrees of freedom are mixed by the anharmonicity. The main consequence of the mixing is that ω_a depends also on the total energy, and not only on the axial energy. Since the axial and radial oscillation periods differ by a factor of about 500, it is possible to define a local axial frequency which varies during the radial motion of the atoms [11]

$$\omega_a(r) = 2\pi\sqrt{2U_0\exp(-2r^2/w^2)/M\lambda_T^2}\,. \tag{0.2}$$

As a result, the discrete excitation frequencies shown in Fig. 3 for the 1D case become bands in 3D. The width of these levels is expected to be negligible for atoms at the bottom of the trap, which experience an harmonic potential, and to increase for increasing total energy, because of the longer time spent by the atoms in the anharmonic part of the potential. Thus, while the energetic atoms can be excited by a large range of modulation frequencies on the red of each parametric resonance, the ones at the bottom of the potential can be excited only by a narrow range of frequencies close to ω_a and its harmonics, as confirmed by the temperature spectrum in Fig. 2.

The role of anharmonicity in the excitation of atoms having different energies is confirmed by a measure of the equilibration time for the radial degree of freedom, following a short axial excitation period, as reported in Fig. 4. When Ω is tuned to the red of $2\,\omega_a$, the equilibration time is of the order of 5 ms, almost an order of magnitude shorter than the elastic collisional time expected for our sample [9]. This observation indicates that anharmonic cross-dimensional mixing can actually be the dominant process in this case, as expected for a perturbation of the high-lying trap states. If Ω is tuned to $2\,\omega_a$ the two degrees of freedom thermalize on a longer timescale of 20-50 ms, which is more likely to be determined by the rate of interatomic collisions alone. In fact the thermalization time shows an almost linear dependence on the atomic density, as already observed in Refs. [12,9]. In accordance with our heuristic arguments, it appears that anharmonic mixing is not important for excitation of atoms at the bottom of the trap.

We have investigated the time evolution of losses and temperature for modulation frequencies red-detuned from $2\,\omega_a$, to get an indication of the efficiency of the selective parametric excitation in cooling the atoms in the FORT. The evolution reported in Fig. 5 was obtained with ϵ=12%, $\Omega = 1.6\,\omega_a$, and T_{ex} variable between 5 and 300 ms; the atom number and temperature were detected after a 20-ms thermalization period. The general features of the process discussed above are thus confirmed: at the beginning of the excitation phase the energetic atoms are very efficiently removed from the trap, and very likely the selection is performed on the total (3D) energy (the reduction of the radial temperature following axial excitation is very fast). We think that elastic collisions are responsible for a continuous replenishing of the high-lying states of the trap, giving rise to the observed exponential decrease of both atom number and temperature on a timescale much longer than 5 ms. Thus we can consider this process to be one of evaporative cooling, forced by the parametric excitation. As shown in Fig. 5, after 150 ms the atom number and temperature reach a steady state value, which is conserved even after a further 150 ms period of excitation. The features just described are very interesting from the perspective of using this technique to cool the atoms in the trap, since an excitation at the proper frequency provides a fast removal of the energetic atoms without heating the coldest ones. After switching-off the parametric modulation at t=300 ms we have measured a temperature increase at a rate of 34(12) μKs^{-1} (compatible with that expected for photon scattering), which indicates that the minimum temperature reached in the cooling phase is probably not limited by the excitation process itself.

From the two sets of data in Fig. 5 we extract the evolution of the relative phase space density ρ reported the inset, assuming thermal equilibrium between radial and axial motions. Our experimental value for the parameter which describes the overall efficiency of an evaporative cooling process [13] is

$$\gamma_{tot} = \frac{N_f \rho_f}{N_i \rho_i} = 0.7(1)\,. \tag{0.3}$$

It can be compared to the efficiency of a variety of evaporative techniques in optical and magnetic traps, as reported in Ref. [13]. To our knowledge the process we have investigated here represents the only way of forcing evaporative cooling in a standard optical trap [14] without a reduction in the trap depth, a technique investigated in Ref. [15].

Our FORT is actually not optimized for evaporation; indeed we are obliged by the short lifetime of the trap to force the removal of atoms on a timescale not much larger than the elastic collisional time. Further investigations in a better vacuum environment, or at a larger atomic density, can provide better tests of the efficiency of this cooling technique. Indeed, with a longer time for evaporation one could try to use a modulation frequency on the red of all the axial resonances ($\Omega < \omega_a$), to excite only the very energetic atoms and thus increase the cooling efficiency. Also, one could try to investigate the effect of a sweep of Ω towards resonance during cooling, in an analogous way to what is usually performed in magnetic traps.

In conclusion, we have reported the observation of an energy-selective removal of atoms from a FORT, as a consequence of parametric excitation of anharmonic vibrational modes. In addition, we have shown that the energy selectivity can be exploited to cool the trapped sample. Since this effect arises due to the anharmonicity of the trapping potential, it should be observable in most actual traps for cold particles. The removal mechanism should work independently of the internal structure of the trapped particles (atoms, molecules or ions), as it is based on the excitation of their external degrees of freedom. The observations reported here should provide the impetus for further investigations and theoretical modeling necessary for characterizing the ultimate limits and efficiency of this cooling technique; however, it already appears interesting for a fast cooling of an atomic sample in a tight FORT.

We thank M. Inguscio for useful suggestions and for careful reading of the manuscript. We also acknowledge useful discussions with R. Jáuregui, and we thank Scuola Normale Superiore, Pisa for the loan of the trapping laser. This work was supported by the ECC under the Contract HPRICT1999-00111, and by MURST under the PRIN 2000 Program. R.B. was partially supported by ASI.

---

[1] T. A. Savard, K. M. O'Hara, and J. E. Thomas, Phys. Rev. A**56**, R1095 (1997); M. E. Gehm, K. M. O'Hara, T. A. Savard, and J. E. Thomas, Phys. Rev. A **58**, 3914 (1998).

[2] C. W. Gardiner, J. Ye, H. C. Nagerl, and H. J. Kimble, Phys. Rev. A **61**, 045801 (2000).

[3] K. M. O'Hara, S. R. Granade, M. E. Gehm, and T. A. Savard, Phys. Rev. Lett. **82**, 4204 (1999).

[4] H. Engler, T. Weber, M. Mudrich, R. Grimm, and M. Weidemüller, Phys. Rev. A **62**, 031402 (2000).

[5] S. Friebel, C. D'Andrea, J. Walz, M. Weitz, and T. W. Hänsch, Phys. Rev. A **57**, R20 (1998).

[6] V. Vuletic, C. Chin, A. J. Kerman, and S. Chu, Phys. Rev. Lett. **81**, 26(1998).

[7] J. Ye, D. W. Vernooy, and H. J. Kimble, Phys. Rev. Lett. **83**, 4987 (1999).

[8] R. Jáuregui, N. Poli, G. Roati, and G. Modugno, submitted to Phys. Rev. A; arXiv: physics/0103046.

[9] G. Roati, W. Jastrzebski, A. Simoni, G. Modugno, and M. Inguscio, to appear in Phys. Rev. A; arXiv: physics/0010065.

[10] A general result of the theory presented in [1] is that the excitation rate is proportional to $\epsilon^2\omega^2$.

[11] T. Elsässer, diploma thesis, University of Heidelberg (2000).

[12] V. Vuletic, A. J. Kerman, C. Chin, and S. Chu, Phys. Rev. Lett. **82**, 1406 (1999).

[13] W. Ketterle and N. J. van Druten, Adv. At. Mol. Opt. Phys. **37** 181 (1996).

[14] An exception is represented by the optical trap investigated in K.L. Corwin, S. J. M. Kuppens, D. Cho, and C. E. Wieman, Phys. Rev. Lett. **83**, 1311 (1999).

[15] C. S. Adams, H. J. Lee, N. Davidson, M. Kasevich, and S. Chu, Phys. Rev. Lett. **74**, 3577 (1995).



FIG. 1. Spectrum of the resonances in the trap losses (open circles) and in the radial temperature (solid circles) following a T_{ex}=50 ms, ϵ=30% excitation of the radial vibration. The shaded area indicates the initial temperature range, and represents the uncertainty in temperature.



FIG. 2. Spectrum of the resonances in the trap losses (open circles) and in the radial temperature (solid circles) following a T_{ex}=200 ms, ϵ=16% excitation of the axial vibration.



FIG. 3. Calculated spectrum of the axial excitation frequency for a 1D optical lattice with U_0=700 μK.



FIG. 4. Equilibration of the radial degree of freedom after an excitation at 1.8ω_a (triangles), with T_{ex}=2 ms and ϵ=12%, and at 2ω_a (circles), with T_{ex}=10 ms and ϵ=12%. The solid curves represent the best fit with an exponential equilibration, plus a constant heating rate for the lower set of data.



FIG. 5. Decrease of the radial temperature (triangles) and of the number of atoms (circles), during a parametric axial excitation with ϵ=12% and Ω=1.6 ω_a. The increase in relative phase-space density is shown in the inset